

YOUR TREATMENT GUIDE



EXPERIENCE THE CALM

CONTROL STEQEYMA CAN BRING

Get to know STEQEYMA, an FDA-approved biosimilar to Stelara[®] (ustekinumab)

STEQEYMA is a prescription medicine used to treat:

- **Moderate to severe plaque psoriasis** in adults and children 6 years and older who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet light alone or with pills)
- **Active psoriatic arthritis** in adults and children 6 years and older
- **Moderately to severely active Crohn's disease** in adults
- **Moderately to severely active ulcerative colitis** in adults

It is not known if STEQEYMA is safe and effective in children less than 6 years of age.

Do not take STEQEYMA if you are allergic to ustekinumab products or any of the ingredients in STEQEYMA.

IMPORTANT SAFETY INFORMATION

Serious infections. STEQEYMA affects your immune system and can increase the risk of serious infections, including tuberculosis (TB), bacterial, fungal, and viral infections. Some infections may require hospitalization.

FDA, Food and Drug Administration.

Models are used for illustrative purposes only. Not actual patients.



Please see Important Safety Information throughout and on pages 12-13. Full Prescribing Information, including Medication Guide, is available [here](#).

What is STEQEYMA?

STEQEYMA is a prescription biosimilar

STEQEYMA is a biosimilar of Stelara®. Biosimilar means that:

- The biological product is approved based on data demonstrating that it is highly similar to an FDA-approved biological product, known as a reference product
- There are no clinically meaningful differences in how it works or in how safe it is compared to the reference product

A biosimilar is typically more affordable than the reference product.

Since STEQEYMA is a biosimilar of Stelara, it can be used instead of Stelara.

Similar to Stelara, STEQEYMA works on the immune system to protect your body from inflammation

STEQEYMA shuts down signals in your immune system, preventing them from activating certain inflammatory immune cells. By doing so, STEQEYMA can reduce inflammation.




IMPORTANT SAFETY INFORMATION continued

Before starting STEQEYMA:

- Your doctor will test you for TB.
- If you're at risk for TB, you may need treatment before and during STEQEYMA.

Why did my doctor prescribe STEQEYMA?

Your doctor prescribed STEQEYMA to treat your immune-related condition. STEQEYMA has been shown to help:



Moderate to severe plaque psoriasis in adults and children 6 years and older who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet light alone or with pills)

Plaque psoriasis is a chronic autoimmune skin condition characterized by the formation of raised red and scaly patches on the skin. These patches, or plaques, often have a silvery-white appearance, can be itchy and painful, and can sometimes bleed.




Active psoriatic arthritis in adults and children 6 years and older

Psoriatic arthritis is a type of inflammatory arthritis affecting some people with plaque psoriasis. It can affect joints on one or both sides of the body, causing pain, swelling, and stiffness. The joints most commonly affected are the fingers and toes, but it can also affect other joints such as the knees, hips, and spine.



Moderately to severely active Crohn's disease in adults

Crohn's disease is a type of inflammatory bowel disease that causes chronic inflammation in the digestive tract. It can affect any part of the digestive tract, from the mouth to the anus.



Moderately to severely active ulcerative colitis in adults

Ulcerative colitis is a chronic inflammatory bowel disease that affects the colon and rectum. It is characterized by inflammation and ulcers (sores) in the lining of the colon and rectum, which can lead to symptoms such as abdominal pain, diarrhea, and rectal bleeding.

IMPORTANT SAFETY INFORMATION continued


• Tell your doctor if you:

- Have an infection or signs of infection: fever, chills, cough, muscle aches, shortness of breath, weight loss, blood in phlegm, tiredness, diarrhea, stomach pain, skin sores, or burning when urinating.

How STEQEYMA can help

STEQEYMA was shown to be as effective and safe as Stelara®

In a clinical study of 509 patients with moderate to severe plaque psoriasis, STEQEYMA was shown to be as effective as Stelara in reducing the severity of symptoms and decreasing the area affected.



STEQEYMA and Stelara delivered similar results—**67%** of STEQEYMA patients and **61%** of Stelara patients achieved **90%** improvement in plaque psoriasis symptoms.

Individual results may vary. See the [Medication Guide](#) for full uses for STEQEYMA.

IMPORTANT SAFETY INFORMATION continued

- Are being treated for an infection or have any open cuts.
- Get frequent or recurring infections.
- Have TB or have been exposed to TB.
- Ever had an allergic reaction to ustekinumab products.



STEQEYMA is approved to treat other diseases it has not been directly tested on in studies. The FDA looks at a wide variety of tests and data to determine that a drug is biosimilar to a reference product, including clinical trials, blood and immune-system evaluations, and chemical analyses. STEQEYMA has been approved by FDA as a biosimilar to Stelara.



FDA, Food and Drug Administration.

IMPORTANT SAFETY INFORMATION continued

- Recently had or plan to get vaccines (you must avoid live vaccines, including the BCG vaccine during treatment and for a year afterward).
- Have new or changing lesions within plaque psoriasis areas or on normal skin.

Side effects for STEQEYMA

Common side effects of STEQEYMA include:

- nasal congestion, sore throat, and runny nose
- upper respiratory infections
- fever
- headache
- tiredness
- itching
- nausea and vomiting
- redness at the injection site
- vaginal yeast infections
- urinary tract infections
- sinus infection
- bronchitis
- diarrhea
- stomach pain

These are not all of the possible side effects of STEQEYMA.

You should also review **“What is the most important information I should know about STEQEYMA?”** in the [Medication Guide](#) or on pages 12 and 13 of this brochure.



Call your doctor for medical advice about side effects.
You may report side effects to FDA at 1-800-FDA-1088.


You may also report side effects to Celltrion USA, Inc.
at 1-800-560-9414.

Designed for safe and efficient administration


STEQEYMA is available in single-dose, prefilled syringe injections that offer you the flexibility and control of self-administration

You'll be shown how to use the injection by your doctor or a specialized nurse so you can give yourself the injection at home.

STEQEYMA prefilled syringes are available in 2 dose strengths




45-mg/0.5-mL, single-dose PFS



90-mg/1.0-mL, single-dose PFS

The STEQEYMA passive needle guard prevents needle-stick injuries, for added safety during administration. Plus, its ergonomic design allows easy, one-handed injection activation, for comfort and convenience during use.

Parts of the prefilled syringe injection



Do not try to inject STEQEYMA yourself until you or your caregiver have been shown how to inject STEQEYMA by your doctor or nurse.

PFS, prefilled syringe.

IMPORTANT SAFETY INFORMATION continued

- Are receiving or have received allergy shots or phototherapy.
- Are pregnant, planning to become pregnant, or breastfeeding. STEQEYMA can pass into breast milk.
- Received STEQEYMA while pregnant, inform your baby's healthcare provider before they get vaccinations.

Watch the self-injection video [here](#).

IMPORTANT SAFETY INFORMATION continued

- **Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know all the medicines, vitamins, and supplements you take, and share them with your doctor.

Designed for safe and efficient administration (continued)

Convenient administration



Finger flange to improve grip comfort

The inclusion of a **finger flange** in the injection design aims to enhance grip comfort by providing an ergonomic extension for the fingers, allowing for a more secure and comfortable grip on the device.



Storage for up to 15 days at room temperature (up to 86°F)

STEQEYMA demonstrates stability for a duration of up to **15 days** when stored at room temperature, specifically at a **maximum of 86°F**.

At room temperature, the syringe must be kept in the original carton to protect it from light and discarded if not used within the 15-day period.


STEQEYMA prefilled syringes can be stored up to 3 years in a refrigerator (36° to 46°F).

IMPORTANT SAFETY INFORMATION continued

After starting STEQEYMA:

Call your doctor right away if you develop symptoms of an infection, such as chest infections, skin infections, or shingles. STEQEYMA can increase the risk of infections or make existing infections worse.

Less injection-related pain, allergy, and injury



A 27-gauge special thin wall (STW) needle

A **27-gauge STW needle** is employed in the STEQEYMA device to alleviate needle-related anxiety and pain perception.

How thin is the STEQEYMA needle?

Regular Wall



Thin Wall



Special Thin Wall




Removal of citrate buffers

By being **citrate free**, the injection is designed to reduce discomfort associated with self-injections, as citrate buffers are known to cause pain.



Does not contain natural rubber latex

The STEQEYMA device is designed to be **latex free**, ensuring that patients with latex allergies can safely utilize it without any risk of adverse reactions.



Passive needle guard

By incorporating a **passive needle guard**, the STEQEYMA device provides a protective measure to prevent any accidental needle stick injuries during use.

IMPORTANT SAFETY INFORMATION continued

Cancers

STEQEYMA may increase the risk of certain cancers, including skin cancer. Tell your doctor if you have a history of cancer or notice any new skin growths during treatment.

IMPORTANT SAFETY INFORMATION

Serious infections. STEQEYMA affects your immune system and can increase the risk of serious infections, including tuberculosis (TB), bacterial, fungal, and viral infections. Some infections may require hospitalization.

Before starting STEQEYMA:

- Your doctor will test you for TB.
- If you're at risk for TB, you may need treatment before and during STEQEYMA.
- **Tell your doctor if you:**
 - Have an infection or signs of infection: fever, chills, cough, muscle aches, shortness of breath, weight loss, blood in phlegm, tiredness, diarrhea, stomach pain, skin sores, or burning when urinating.
 - Are being treated for an infection or have any open cuts.
 - Get frequent or recurring infections.
 - Have TB or have been exposed to TB.
 - Ever had an allergic reaction to ustekinumab products.
 - Recently had or plan to get vaccines (you must avoid live vaccines, including the BCG vaccine during treatment and for a year afterward).
 - Have new or changing lesions within plaque psoriasis areas or on normal skin.
 - Are receiving or have received allergy shots or phototherapy.
 - Are pregnant, planning to become pregnant, or breastfeeding. STEQEYMA can pass into breast milk.
 - Received STEQEYMA while pregnant, inform your baby's healthcare provider before they get vaccinations.
- **Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know all the medicines, vitamins, and supplements you take, and share them with your doctor.

After starting STEQEYMA:

Call your doctor right away if you develop symptoms of an infection, such as chest infections, skin infections, or shingles. STEQEYMA can increase the risk of infections or make existing infections worse.

Cancers

STEQEYMA may increase the risk of certain cancers, including skin cancer. Tell your doctor if you have a history of cancer or notice any new skin growths during treatment.

Posterior Reversible Encephalopathy Syndrome (PRES)

PRES is a rare but serious brain condition that can cause death. Symptoms include headache, confusion, seizures, and vision problems. Contact your doctor immediately if these occur.

Serious Allergic Reactions. Serious allergic reactions can occur with STEQEYMA. Stop using STEQEYMA and get medical help right away if you have any of the following symptoms of a serious allergic reaction: feeling faint, chest tightness, swelling of your face, eyelids, tongue, or throat, or a skin rash.

Lung Inflammation. Cases of lung inflammation have happened in some people who receive ustekinumab products and may be serious. These lung problems may need to be treated in a hospital. Tell your doctor right away if you develop shortness of breath or a cough that doesn't go away during treatment with STEQEYMA.

Common Side Effects of STEQEYMA. Common side effects of STEQEYMA include nasal congestion, sore throat, and runny nose; upper respiratory infections; fever; headache; tiredness; itching; nausea and vomiting; redness at the injection site; vaginal yeast infections; urinary tract infections; sinus infection; bronchitis; diarrhea; and stomach pain.

These are not all of the possible side effects of STEQEYMA. Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Celltrion USA, Inc. at 1-800-560-9414.

What should I know before self-injecting?

Make sure you speak with your doctor

You should consult with your doctor before you start your STEQEYMA injections. Use STEQEYMA exactly as your doctor tells you to.

It is also important that you know the do's and don'ts of handling the device.

DO

Leave syringe outside of the carton for 30 minutes to allow it to reach room temperature.

Check the expiration date on the carton and the label of the prefilled syringe.

Make sure you have the correct medicine and dosage.

Check the prefilled syringe to make sure it is not cracked or damaged.

Look at the medicine and confirm that the liquid is clear to slightly opalescent and colorless to pale yellow.

DON'T

Do not warm the prefilled syringe using heat sources such as hot water or a microwave.

Do not use the injection if the expiration date has passed.

Do not use the injection if you are allergic to ustekinumab or any of the other ingredients of STEQEYMA.

Do not use the injection if you have an active infection which your doctor thinks is important.

Do not shake the prefilled syringes or use the injection if the medicine has been shaken strongly.



IMPORTANT SAFETY INFORMATION continued

Posterior Reversible Encephalopathy Syndrome (PRES)

PRES is a rare but serious brain condition that can cause death. Symptoms include headache, confusion, seizures, and vision problems. Contact your doctor immediately if these occur.

IMPORTANT SAFETY INFORMATION continued

Serious Allergic Reactions. Serious allergic reactions can occur with STEQEYMA. Stop using STEQEYMA and get medical help right away if you have any of the following symptoms of a serious allergic reaction: feeling faint, chest tightness, swelling of your face, eyelids, tongue, or throat, or a skin rash.

Patient resources available for you

These tools can help you manage and make the most of your treatment



Self-injection video

Learn how to self-inject by watching this brief video with clear instructions.



Self-injection brochure/calendar

Use this helpful tool to keep track of your injections and treatment schedule.

View complete resources [here](#).

IMPORTANT SAFETY INFORMATION continued

Lung Inflammation. Cases of lung inflammation have happened in some people who receive ustekinumab products and may be serious. These lung problems may need to be treated in a hospital. Tell your doctor right away if you develop shortness of breath or a cough that doesn't go away during treatment with STEQEYMA.

Celltrion support programs

These programs help you get access to STEQEYMA



Help verify your insurance coverage for STEQEYMA, determine your plan-specific requirements, and understand your available benefits.



Connect with Celltrion CARES® for co-pay assistance with your commercial insurance.



Helps you get started with your STEQEYMA treatment through expert nurse-supported injection training, storage and handling guidance, dosing education, and ongoing support.*

*While Nurse Connectors are registered healthcare professionals, they cannot offer medical advice. If you have an urgent or specific question related to your healthcare, please contact your HCP.



IMPORTANT SAFETY INFORMATION continued

Common Side Effects of STEQEYMA. Common side effects of STEQEYMA include nasal congestion, sore throat, and runny nose; upper respiratory infections; fever; headache; tiredness; itching; nausea and vomiting; redness at the injection site; vaginal yeast infections; urinary tract infections; sinus infection; bronchitis; diarrhea; and stomach pain.

YOUR TREATMENT GUIDE

 **SteQeyma**[®]
(ustekinumab-stba)

Watch the
self-injection
video [here](#).

Please see Important Safety Information throughout and on pages 12-13.
Full Prescribing Information, including Medication Guide, is available [here](#).

Stelara is a registered trademark of Johnson and Johnson and is marketed by Janssen Biotech, Inc. STEQEYMA is a registered trademark of Celltrion, Inc., used under license. Celltrion CARES is a registered trademark of Celltrion, Inc., used under license. Celltrion CONNECT is a registered trademark of Celltrion Holdings Co., Ltd., used under license.

©Celltrion USA, Inc. 2025. US-STEQ-25-00019.

 **CELLTRION** USA